

ACTIVE PHYTO & MULTI COMPLEX



Clinical Applications

- Foundational Nutrition*
- Basic Formula for Wellness*
- Supports Antioxidant Activity*
- Supports Detoxification*
- Supports Health in Individuals with Inadequate Nutrient Intake*
- Supports Energy Production and Stress Response*

*Active Phyto & Multi Complex is a premium multivitamin/mineral formula with metabolically active B vitamins and a bioavailable mineral complex for optimal nutrient utilization. In addition to foundational nutrition, this comprehensive formulation features powerful phytochemical sources, such as astaxanthin and extracts of green coffee bean, green tea, and rosemary, to support the body's antioxidant and cell-protective mechanisms and to counter oxidative stress.**

All Pharmacist Formulations Formulas Meet or Exceed cGMP Quality Standards

Discussion

Active Phyto & Multi Complex provides enhanced antioxidant support through a special phytonutrient blend. The primary mechanisms of action of each ingredient unique to Active Phyto & Multi Complex are described below, followed by a discussion of the multivitamin and mineral components that provide the foundation for all the Active Phyto & Multi Complex formulations.*

- **Astaxanthin** is a naturally occurring xanthophyll carotenoid with a distinctive molecular structure that allows it to scavenge and quench reactive oxygen species and free radicals in the inner and outer layers of the cell membrane.¹ Astaxanthin has been studied extensively for systemic benefits because of its strong antioxidant activity, which research has indicated is much greater than vitamins C and E, coenzyme Q10, green tea catechins, lutein, or other carotenoids.² Astaxanthin is receiving increasing attention as an effective molecule to prevent oxidative stress-mediated and age-related decline.*^{1,2}
- **CinSulin®** is a patented water extract of cinnamon. The unique proprietary extraction and dehydration process for manufacturing CinSulin results in a concentrated (10:1) extract that minimizes undesirable substances while retaining those that are health-promoting, such as type-A polyphenolic polymers.³ Cinnamon has been used for hundreds of years as a flavor additive and in traditional Eastern medicine for its health-promoting qualities.⁴ Cinnamon bark has high antioxidant activities due to the presence of polyphenols and volatile oil compounds, and cinnamon has been studied extensively for its roles in glucose uptake, glycogen synthesis, insulin action, and support for healthy blood lipid metabolism.*⁴⁻⁶
- **Green Coffee Bean** extract contains polyphenols with strong antioxidant and free radical scavenging qualities that are associated with reducing oxidative stress and its complications.^{7,8} The major polyphenols from green coffee beans are collectively known as chlorogenic acids; these components or their metabolites mediate many benefits associated with green coffee bean extract.⁸ The effects of these phytonutrients on vascular health, body composition, and glycemic and lipid profiles are areas of interest for human research.*^{8,9}
- **Maca Root** has been used as a food and traditional medicine in the Andes of Peru for over 2,000 years.¹⁰ Both classical practices and modern science endorse maca's vitality-, aphrodisiac-, and fertility-promoting qualities.¹⁰ Characteristic of many folk medicines, maca has numerous applications. In animal studies, maca demonstrates antioxidant, neuroprotective, and antiviral activities.¹¹ Research attributes the benefits of maca root to its antioxidant compounds, such as phenols, sterols, glucosinolates, alkamides, and polysaccharides, as well as to its secondary metabolites, including macaene and macamide.*^{10,11}
- **Green Tea** leaves are a source of polyphenols known as catechins, with epigallocatechin-3-gallate (EGCG) being the most abundant and biologically active catechin in green tea leaves. Green tea extracts are considered more stable than pure EGCG because of the broad spectrum of antioxidant constituents in the extract.¹² In vitro research demonstrates that green tea polyphenols have direct antioxidant activity by scavenging reactive oxygen species or chelating transition metals.¹³ Green tea polyphenols may also act indirectly by upregulating phase II antioxidant enzymes.¹³ In vivo research shows that these polyphenols increase plasma antioxidant activity, suppress oxidative stress markers, and protect against degenerative processes.*^{12,13}
- **Rhodiola** is an adaptogenic herb used traditionally in Eastern Europe and Asia for centuries. Medical and pharmacological texts describe its use for immune, psychiatric, and neurological health. In healthy individuals, therapeutic doses have helped relieve fatigue and increase attention span, memory, and work productivity.¹⁴ Rhodiola root is rich in salidroside and rosavin, phenolic compounds that have strong antioxidant properties. Research into Rhodiola's antioxidant activity has revealed its protective effects against oxidative damage in the nervous system and how it impacts memory and cognition by improving resistance to physical and emotional stress.*^{14,15}
- **Milk Thistle** seeds have been used for medicinal purposes for over 2,000 years. Silymarin, which consists of a mixture of flavonolignans and the flavonoid taxifolin, is the main active component of milk thistle.¹⁶ Silymarin's primary site of action is in the liver, where it can play a role in maintaining healthy levels of fat peroxidation, moderating fibrous tissue formation, supporting healthy immune and

Continued on next page

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**



Distributed By:
Pharmacist Formulations, LLC
6506 North Davis Highway
Pensacola, FL 32504
www.PharmacistFormulations.shop

ACTIVE PHYTO & MULTI COMPLEX



inflammatory responses, and promoting protein synthesis and normal liver tissue regeneration.¹⁶ Silymarin supports antioxidant activity, neutralizes toxins, and may have hepatoprotective effects.^{*16-18}

- **Rosemary** is an aromatic plant originating from the Mediterranean region and is the most common culinary herb cultivated worldwide. It is rich in phytochemicals with antioxidant attributes and cytokine-balancing properties.^{19,20} The antioxidant properties of rosemary extract are well-established and primarily attributed to phenolic compounds, including diterpenes, triterpenes, carnosic acid, ursolic acid, and rosmarinic acid. Carnosic and rosmarinic acids demonstrate the most prevalent physiological effects and interact with multiple molecular targets.^{*19,20}
- **Bitter Melon** has a rich history of use in traditional and folk medicine. Its fruit has a unique phytochemistry with bioactive components, including terpenoids, glycosides, flavonoids, alkaloids, tannins, and numerous saponins. These components are responsible for a wide range of activities, including antioxidant, cytokine balancing, free radical scavenging, and moderating NF- κ B and TNF- α cascades.^{21,22} These mechanisms confer protection to cellular and organ function.²¹
- **Artichoke** has been cited as a medicine since ancient times, primarily regarding its use for digestive and liver health.^{23,24} Modern research has validated these historic uses.²³ Polyphenolic compounds, such as cynarin, chlorogenic acid, and flavonoids, have been documented as artichoke's active principles and are present mainly in the leaves. In vitro research demonstrated that artichoke leaf extract's antioxidant potential is due to the radical scavenging and metal ion-chelating effects of its unique blend of polyphenol constituents.^{*23}

Active Phyto & Multi Complex—Foundation Formula Discussion

Adequate nourishment is the foundation for overall health and wellness, and good nutrition typically translates into a stronger immune system and better health. The human body uses dietary proteins, fats, and carbohydrates, known as macronutrients, to provide the energy (calories) needed to fuel physiological functions. Vitamins and minerals, known as micronutrients, are needed in much smaller quantities. Unlike their macro counterparts, micronutrients don't give you energy, but they do participate in converting food to energy; building and repairing tissues and DNA; manufacturing neurotransmitters, hormones, and other modulators in the body; breaking down and detoxifying xenobiotics and medications; and maintaining growth, reproduction, and health.^{*25-27}

According to the Dietary Guidelines for Americans 2020-2025 (DGA) and additional data from the USDA and other agencies and organizations, the American diet lacks micronutrients.²⁸⁻³⁰ Mass food production, storage techniques, poor food choices, and nutrient-depleting preparation methods may be contributing to this deficit. Furthermore, the percent daily values (%DV) for micronutrients are based on the minimum amount needed to meet the basic need of a healthy person of a specific age and gender group. The %DV is not always indicative of the amount needed for optimal functioning of all individuals, especially those who are chronically ill.^{*27,29,31}

When considering where American diets fall short in nutrients, the DGA shows that low intakes of potassium, dietary fiber, calcium, and vitamin D are a public health concern.²⁸ Other nutrients that have notably low intakes or require increased intake subsequent to life stage include vitamins A, B6, B12, C, E, and folate; the minerals magnesium and iron; and choline.^{28,32,33} Data from the National Health and Nutrition Examination Surveys (NHANES) suggest a pervasive deficiency in A, C, D, E, and zinc—nutrients linked to immune health.³⁰ Inadequate intake of most of these nutrients is attributable to an overall unhealthy eating pattern due to low intakes of nutrient-rich foods such as vegetables, fruits, whole grains, and dairy that contain these nutrients.²⁸ In cases when food is not enough for an individual to get adequate micronutrients, multivitamin/mineral supplements are recognized as being of value to help fill dietary nutritional shortfalls.^{*26,30,31,34-36}

Active Phyto & Multi Complex is designed to meet the foundational nutrition needs for a variety of protocols and life stages. This formula provides:

A Balanced Profile Vitamins and minerals work cooperatively when present in sufficient amounts. However, imbalances between micronutrients can disrupt this synergistic relationship, possibly leading to instances of competitive intestinal absorption or displacement at the metabolic/cellular level, which can produce relative excesses and insufficiencies. For this reason, Active Phyto & Multi Complex features a balanced nutrient profile that includes calcium and magnesium, vitamins C and E, bioactive folate, bioactive vitamin B12, B vitamin complex, beta-carotene, and trace elements.*

Bioavailable Nutrient Forms The micronutrients are provided in bioactive forms so that they can be adequately absorbed and utilized. Active Phyto & Multi Complex contains a full complement of Albion® patented mineral chelates and complexes. Albion is a recognized world leader in mineral amino acid chelate nutrition and manufactures highly bioavailable nutritional mineral forms that are validated by third-party research and clinical studies. Active Phyto & Multi Complex also contains natural vitamin E, clinically shown to be more bioavailable than synthetic dl-alpha-tocopherol, as well as mixed tocopherols to more closely approximate how much vitamin E an individual might gain when consuming healthful foods.^{37,38} The folate source in Active Phyto & Multi Complex is methyltetrahydrofolate (5-MTHF)—the most bioactive form of folate³⁹—in the form of Quatrefolic®, which has greater stability, solubility, and bioavailability over calcium salt forms of 5-MTHF. Supplementing with bioactive 5-MTHF facilitates the bypassing of steps in folate metabolism. This may be especially beneficial to individuals with genetic variations in folate metabolism.^{40,41} Vitamins B2, B6, and B12 are provided in metabolically active forms.*

Support for Energy Production and Stress Response Active Phyto & Multi Complex provides generous levels of B vitamins, which serve as prime coenzymes in glycolysis and oxidative phosphorylation and as cofactors in amino acid and lipid metabolism.⁴²⁻⁴⁴ Sufficient levels of the B vitamins are critical for energy production and cell growth and division, and they have many other essential roles in the body, including support for nervous system function.⁴⁵ The balanced presence of B vitamins is essential to their cooperative functioning and are excellent for individuals with stressful lifestyles.*

Antioxidant Protection Vitamins E and C, selenium, zinc, beta-carotene, and trace elements provide broad-spectrum antioxidant activity.^{46,47} Their combined presence supports their ability to regenerate each other and maintain consistent levels of antioxidant activity both intra- and extracellularly.*

Detoxification Support Xenobiotics, including environmental pollutants and medications, must undergo biotransformation into molecules that can be easily excreted from the body. Detoxification of xenobiotics is a complex process that requires micronutrients, phytonutrients, energy, and adequate antioxidant support for safe and effective completion.⁴⁷⁻⁴⁹ There are significant levels of bioavailable riboflavin, niacin, folate, and B12 present in Active Phyto & Multi Complex to support phase I detoxification. Beta-carotene, vitamin C, tocopherols, selenium, zinc, and manganese are present to support tissues when reactive intermediates are formed between phase I and phase II detoxification.*

Active Phyto & Multi Complex offers foundational multivitamin and mineral support designed to compensate for dietary nutritional shortfalls and nourish optimal wellness. A select combination of ingredients has been added for enhanced support of antioxidant mechanisms.*



Supplement Facts

Serving Size: 2 Capsules
Servings Per Container: 45

Amount Per Serving	%DV	Amount Per Serving	%DV
Vitamin A (as natural beta-carotene and retinyl palmitate)	380 mcg 42%	Chromium (as chromium nicotinate glycinate chelate) ^{S3}	85 mcg 243%
Vitamin C (as sodium ascorbate, potassium ascorbate, zinc ascorbate, and calcium ascorbate)	40 mg 44%	Molybdenum (as molybdenum glycinate chelate) ^{S4}	5 mcg 11%
Vitamin D3 (cholecalciferol)	0.85 mcg (34 IU) 4%	Proprietary Phytonutrient Blend	346.5 mg **
Vitamin E (as d-alpha tocopheryl succinate)	20 mg 133%	Green Coffee Extract (<i>Coffea arabica</i> , <i>Coffea robusta</i>)(seed)(15% total chlorogenic acids), Organic Maca (<i>Lepidium meyenii</i>)(root), Green Tea Aqueous Extract (<i>Camellia sinensis</i>)(leaf)(30% EGCG), Rhodiola Extract (<i>Rhodiola rosea</i>)(root)(3% rosavins), Milk Thistle Extract (<i>Silybum marianum</i>)(seeds)(30% silybins), Rosemary Extract (<i>Rosmarinus officinalis</i>)(leaf)(5% rosmarinic acid), Bitter Gourd Extract (<i>Momordica charantia</i>)(fruit), Artichoke Extract (<i>Cynara scolymus</i>)(leaf)(7% caffeoylquinic acids), Cinnamon 10:1 Aqueous Extract (<i>Cinnamomum cassia</i>)(bark) ^{S4} , Astaxanthin ^{S4}	
Thiamin (as thiamine mononitrate)	3 mg 250%		
Riboflavin (as riboflavin 5'-phosphate sodium)	3 mg 231%		
Niacin (as niacinamide and niacin)	10 mg 63%		
Vitamin B6 (as pyridoxal 5'-phosphate)	3 mg 176%		
Folate (as (6S)-5-methyltetrahydrofolate acid, glucosamine salt) ^{S1}	110 mcg DFE 28%		
Vitamin B12 (as methylcobalamin)	85 mcg 3,542%		
Biotin	170 mcg 567%		
Pantothenic Acid (as d-calcium pantothenate)	30 mg 600%		
Iodine (as potassium iodide)	15 mcg 10%		
Magnesium (as di-magnesium malate) ^{S2}	15 mg 4%		
Zinc (as zinc bisglycinate chelate) ^{S2}	2 mg 18%		
Selenium (as selenium glycinate) ^{S2}	15 mcg 27%		
Manganese (as manganese bisglycinate chelate) ^{S2}	0.05 mg 2%		

Other Ingredients: Capsule (hypromellose and water), hydroxypropyl cellulose, potassium glycinate, di-calcium malate, ascorbyl palmitate, silica, dicalcium phosphate dihydrate, calcium silicate, and choline diglycinate citrate.

S1. Quatrefolic® is a registered trademark of Gnosis S.p.A.

Produced under U.S. patent 7,947,662.

S2. Aibion® is a registered trademark of Balchem Corporation or its subsidiaries.

S3. CinSin® is registered trademark of Tang-An Medical Co., Ltd. U.S. patent 6,200,569.

S4. Zanthin® is a registered trademark of Valensa International.

Directions

Take two capsules daily, or use as directed by your healthcare professional.

Consult your healthcare professional before use. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

Formulated To Exclude

Wheat, gluten, yeast, soy protein, dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

References

1. Sztretye M, Dienes B, Gönczi M, et al. *Oxid Med Cell Longev*. 2019;2019:3849692. doi:10.1155/2019/3849692
2. E kpe L, Inaku K, Ekpe V. *J Mol Pathophys*. 2018;7(1):1-6. doi:10.5455/jmp.20180627120817
3. The difference between Cinsulin and Cinnamon. Cinsulin. Accessed November 23, 2022. <https://www.cinsulin.com/the-difference-between-cinsulin-and-cinnamon/>
4. Błaszczyk N, Rosiak A, Kałużna-Czaplińska J. *Forests*. 2021;12(5):648. doi:10.3390/f12050648
5. Rao PV, Gan SH. *Evid Based Complement Alternat Med*. 2014;2014:642942. doi:10.1155/2014/642942
6. Anderson RA, Zhan Z, Luo R, et al. *J Tradit Complement Med*. 2016;6(4):332-336. doi:10.1016/j.jtcme.2015.03.005
7. Bosso H, Barbalho SM, de Alvares Goulart R, et al. *Crit Rev Food Sci Nutr*. 2021;1-17. doi:10.1080/10408398.2021.1948817
8. Bothiraj KV, Vanitha V. *Int J Res Pharm Sci*. 2020;11(1):233-240. doi:10.26454/ijrps.v11i1.1812
9. Asbaghi O, Sadeghian M, Nasiri M, et al. *Nutr J*. 2020;19(1):71. doi:10.1186/s12937-020-00587-z
10. Wang Y, Wang Y, McNeil B, et al. *Food Res Int*. 2007;40(7):783-792. doi:10.1016/j.foodres.2007.02.005
11. Korkmaz S. Antioxidants in maca (*Lepidium meyenii*) as a supplement in nutrition. In: Emad S, Azzam GM, eds. *Antioxidants in Foods and Its Applications*. IntechOpen; 2018:178.
12. Chacko SM, Thambi PT, Kuttan R, et al. *Chin Med*. 2010;5:13. doi:10.1186/1749-8546-5-13
13. Forester SC, Lambert JD. *Mol Nutr Food Res*. 2011;55(6):844-854. doi:10.1002/mnfr.201000641.
14. Brown R, Gerbarg P, Ramazanov Z. *HerbalGram*. 2002;56:40-52.
15. Panossian A, Wikman G, Sarris J. *Phytomedicine*. 2010;17(7):481-493. doi:10.1016/j.phymed.2010.02.002
16. Valková V, Ďuranová H, Bilčíková J, et al. *J Microbiol Biotechnol Food Sci*. 2021;2021:836-843.
17. Milk Thistle Fruit. American Botanical Council. Accessed December 27, 2022. <https://www.herbalgram.org/resources/expanded-commission-e/milk-thistle-fruit/>
18. Soleimani V, Delghandi PS, Moallem SA, et al. *Phytother Res*. 2019;33(6):1627-1638. doi:10.1002/ptr.6361
19. Hussain SM, Syeda AF, Alshammari M, et al. *Braz J Med Biol Res*. 2022;55:e11593. doi:10.1590/1414-431X2021e11593
20. Veenstra JP, Johnson JJ. *Int J Nutr*. 2021;6(4):1-10.
21. Bortolotti M, Mercatelli D, Polito L. *Front Pharmacol*. 2019;10:486. doi:10.3389/fphar.2019.00486
22. Saeed F, Alzaal M, Niaz B, et al. *Int J Food Prop*. 2018;21(1):1270-1290. doi:10.1080/10942912.2018.1446023
23. Ben Salem M, Affes H, Ksouda K, et al. *Plant Foods Hum Nutr*. 2015;70(4):441-453. doi:10.1007/s11130-015-0503-8
24. de Falco B, Incerti G, Amato M, et al. *Phytochem Rev*. 2015;14(6):993-1018. doi:10.1007/s11101-015-9428-y
25. Ames BN. *Arch Biochem Biophys*. 2004;423(1):227-234. doi:10.1016/j.abb.2003.11.002
26. Block G, Jensen CD, Norkus EP, et al. *Nutr J*. 2007;6:30. doi:10.1186/1475-2891-6-30
27. Fletcher RH, Fairfield KM. *JAMA*. 2002;287(23):3127-3129. doi:10.1001/jama.287.23.3127
28. Dietary guidelines for Americans, 2020-2025. 9th ed. U.S. Department of Agriculture and U.S. Department of Health and Human Services. December 2020. Accessed December 21, 2022. https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary_Guidelines_for_Americans_2020-2025.pdf
29. Blumberg JB, Bailey RL, Sesso HD, et al. *Nutrients*. 2018;10(2):248. doi:10.3390/nu10020248
30. Reid CA, Chung RY, Devarshi PP, et al. *Nutrients*. 2020;12(6):1735. doi:10.3390/nu12061735
31. Multivitamin/mineral supplements fact sheet. National Institutes of Health. October 12, 2021. Accessed December 12, 2022. <https://ods.od.nih.gov/factsheets/MVMS-HealthProfessional/?print=1>
32. Bird JK, Murphy RA, Ciappio ED, et al. *Nutrients*. 2017;9(7):655. doi:10.3390/nu9070655
33. Multivitamin/mineral (MVM) inclusion in the supplemental nutrition assistance program (SNAP). Council for Responsible Nutrition. Accessed December 13, 2022. <https://www.crnusa.org/multivitamin-mineral-mvm-inclusion-supplemental-nutrition-assistance-program-snap>
34. Blumberg JB, Frei BB, Fulgoni VL, et al. *Nutrients*. 2017;9(8):849. doi:10.3390/nu9080849
35. Blumberg JB, Cena H, Barr SI, et al. *Clin Ther*. 2018;40(4):640-657. doi:10.1016/j.clinthera.2018.02.014
36. Marra MV, Bailey RL. *J Acad Nutr Diet*. 2018;118(11):2162-2173. doi:10.1016/j.jand.2018.07.022
37. Kiyose C, Muramatsu R, Kameyama Y, et al. *Am J Clin Nutr*. 1997;65(3):785-789. doi:10.1093/ajcn/65.3.785
38. Burton GW, Traber MG, Acuff RV, et al. *Am J Clin Nutr*. 1998;67(4):669-684. doi:10.1093/ajcn/67.4.669
39. Venn BJ, Green TJ, Moser R, et al. *Am J Clin Nutr*. 2003;77(3):658-662. doi:10.1093/ajcn/77.3.658
40. Prinz-Langenohl R, Brämwig S, Tobolski O, et al. *Br J Pharmacol*. 2009;158(8):2014-2021. doi:10.1111/j.1476-5381.2009.00492.x
41. Lamers Y, Prinz-Langenohl R, Brämwig S, et al. *Am J Clin Nutr*. 2006;84(1):156-161. doi:10.1093/ajcn/84.1.156
42. Calderón-Ospina CA, Nava-Mesa MO. *CNS Neurosci Ther*. 2020;26(1):5-13. doi:10.1111/cns.13207
43. Kennedy DO. *Nutrients*. 2016;8(2):68. doi:10.3390/nu8020068
44. Depeint F, Bruce WR, Shangari N, et al. *Chem Biol Interact*. 2006;163(1-2):94-112. doi:10.1016/j.cbi.2006.04.014
45. B vitamins. National Library of Medicine. MedlinePlus. September 23, 2021. Accessed December 2, 2021. <https://medlineplus.gov/bvitamins.html>
46. Jayedi A, Rashidi-Pour A, Parohan M, et al. *Adv Nutr*. 2018;1(9):701-716. doi:10.1093/advances/nmy040
47. Doyle ME, Pariza MW. Antioxidant nutrients and protection from free radicals. In: Kotsonis FN, Mackey MA, eds. *Nutritional Toxicology*. 2nd ed. Taylor & Francis; 2002:1-30.
48. Liska DJ. *Altern Med Rev*. 1998;3(3):187-98.
49. Hodges RE, Minich DM. *J Nutr Metab*. 2015;2015:760689. doi:10.1155/2015/760689

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Distributed By:
Pharmacist Formulations, LLC
6506 North Davis Highway
Pensacola, FL 32504
www.PharmacistFormulations.shop

DRS-350
REV. 04/17/23